



## Effect of hot water treatment and oxalic acid at different regimes on storage quality of litchi fruit cv. Rose Scented under ambient conditions

E.S. Marboh\*, R.L. Lal and D.S. Mishra

Department of Horticulture, College of Agriculture, G.B. Pant University of Agriculture & Technology, Pantnagar 263 415

### ABSTRACT

A study was conducted in order to examine the efficacy of hot water treatment (HWT) at 98°C for 30 seconds and 55°C for 12 min. and its combination with oxalic acid (OA) at 0.25 and 10% for 15 min. at room temperature. The control fruits were kept untreated under the same conditions. The fruits were analyzed for physico-chemical characteristics at every one day interval. Combined treatments of HWT (55°C, 12 min) and OA (10%, 15 min.) followed by HWT (98°C, 30 sec) and OA (10%, 15 min.) recorded significantly the lowest level of browning and spoilage loss, and have registered highest marketability percentage, TSS, ascorbic acid content and titratable acidity. However, results suggest that treatment with HWT (55°C, 12 min.) + OA (10%, 15 min.) was most effective in preserving the physiological changes and enhancing quality of litchi fruit.

**Key words:** Litchi, browning index, hot water treatment, oxalic acid, ambient storage.

### INTRODUCTION

Physiological browning and pathological diseases are the two major constraints that limit the marketing scope of litchi in any litchi industry (Zauberman *et al.*, 18). Litchi fruit is very delicate in nature and highly perishable, which accounts for its low shelf life. Wherever it is grown, its shelf life under ambient conditions is never more than 24 to 72 h (Kumar and Kumar, 4). The attractive bright red colour may be lost within 48 hrs (Underhill and Critchley, 16). Studies carried out by workers around the world, have pointed out that pericarp browning in litchi is due to the involvement of several factors *viz.*, enzymatic activities, desiccation, pH (Underhill and Critchley, 17), temperature (chilling injury), mechanical injury, pathogen and pests attack and senescence (Fitzell and Coates, 3).

Sulphur-dioxide fumigation has been used commercially to control pericarp browning (Zauberman *et al.*, 18) but this practice leaves undesirable residues, alters the fruit taste and result in health hazards for consumers and pack house workers (Sivakumar *et al.*, 14). Therefore, maintenance of litchi fruit quality during storage has necessitated the development of an alternative postharvest technology. In this regard, an

investigation was conducted to study the effect of post-harvest treatments, using oxalic acid and hot water treatment at different regimes. Heat treatments have been used to extend storability of several fruits including pomegranate (Mirdehghan *et al.*, 8) and litchi (Lichter *et al.*, 6). Zheng *et al.* (19) reported that use of oxalic acid could give good control of postharvest deterioration mango fruit. The objective of this study was to investigate the effect of various treatments in inhibiting browning and deterioration of litchi fruit during ambient storage.

### MATERIALS AND METHODS

Mature fruit of litchi (*Litchi chinensis* Sonn.) cv. Rose Scented were obtained from an orchard of Horticulture Research Centre, Patharchatta, Uttarakhand. Fruits were selected for uniformity of shape, colour and size and any diseased or blemished fruit discarded. Fruits were destalked (up to 0.5 cm. long pedicels), pre-cooled and subjected to various treatments within 3 hour of harvest. Fruits were subjected to the following different treatments *viz.*, T<sub>1</sub> = Precooling; T<sub>2</sub> = Without Precooling; T<sub>3</sub> = HWT (55°C, 12 min); T<sub>4</sub> = HWT (98°C, 30 sec); T<sub>5</sub> = HWT (55°C, 12 min.) + 0.25% OA (15min.); T<sub>6</sub> = HWT (55°C, 12 min) + 10.0% OA (15min.); T<sub>7</sub> = HWT (98°C, 30 sec) + 0.25% OA (15 min.); T<sub>8</sub> = HWT (98°C, 30 sec) + 10.0% OA (15min.).

Control fruits (precooling, without precooling and HWT) was included for comparison. Precooling was done

\*Corresponding author's present address: Division of Fruits and Horticultural Technology, IARI, New Delhi 110 0012; E-mail: evening\_stone@yahoo.com

**Table 1.** Physiological loss in weight (%), browning index and spoilage percentage of litchi fruits during room temperature storage as affected by various treatments.

Treatment	PLW (%)				Browning index				Spoilage (%)					
	1	2	3	4	1	2	3	4	1	2	3	4	Mean	
T <sub>1</sub>	1.59	2.25	3.27	4.90	36.67	46.67	113.33	210.00	101.67	0.00	11.67	35.00	48.33	23.75
T <sub>2</sub>	1.73	2.51	3.31	5.67	43.33	53.33	133.33	223.33	113.33	0.00	8.33	30.00	43.33	20.42
T <sub>3</sub>	0.87	1.89	3.00	4.53	276.67	296.67	436.67	500.00	377.50	0.00	6.67	26.67	40.00	18.33
T <sub>4</sub>	2.21	2.63	3.83	7.56	366.67	443.33	473.33	500.00	445.83	0.00	8.33	28.33	45.00	20.42
T <sub>5</sub>	1.79	2.56	3.67	5.84	106.67	196.67	266.67	313.33	220.83	0.00	6.67	25.00	35.00	16.67
T <sub>6</sub>	1.89	2.58	3.76	6.56	0.00	0.00	31.67	53.33	21.25	0.00	0.00	11.67	23.33	8.75
T <sub>7</sub>	1.92	3.10	4.69	6.92	56.67	256.67	453.33	500.00	316.67	0.00	6.67	26.67	38.33	17.92
T <sub>8</sub>	2.29	3.17	5.65	9.23	0.00	0.00	32.67	56.67	22.33	0.00	3.33	13.33	26.67	10.83
Means	1.79	2.59	3.90	6.40	110.83	161.67	242.63	294.58	202.43	0.00	6.46	24.58	37.50	17.14
			CD at 5%				CD at 5%					CD at 5%		
Treatment (A)			0.24				10.29					4.12		
Storage days (B)			0.17				7.28					2.91		
A × B			0.49				20.59					8.24		

**Table 2.** TSS, titratable acidity and ascorbic acid content of litchi fruits during room temperature storage as affected by various treatments.

Treatment	TSS (%)				Acidity (%)				Ascorbic acid (mg/ 100 g pulp)										
	0	1	2	3	4	0	1	2	3	4	0	1	2	3	4	Mean			
T <sub>1</sub>	15.93	16.23	16.77	17.37	17.77	16.81	0.46	0.42	0.39	0.36	0.33	0.39	28.91	23.97	19.14	13.35	9.74	19.02	
T <sub>2</sub>	16.03	16.37	16.85	17.53	17.90	16.94	0.45	0.40	0.39	0.35	0.33	0.38	28.86	24.58	17.47	13.81	9.52	18.85	
T <sub>3</sub>	16.15	16.53	16.97	17.53	17.87	17.01	0.41	0.37	0.33	0.32	0.30	0.35	29.02	25.41	18.20	13.81	10.21	19.33	
T <sub>4</sub>	16.09	16.33	16.77	17.63	18.03	16.97	0.47	0.41	0.39	0.34	0.32	0.39	29.30	24.80	17.81	13.81	9.60	19.06	
T <sub>5</sub>	16.23	16.62	17.10	17.43	17.93	17.06	0.43	0.39	0.35	0.32	0.30	0.36	28.58	24.58	19.27	14.24	10.32	19.40	
T <sub>6</sub>	16.23	16.72	17.27	17.83	18.33	17.28	0.45	0.41	0.38	0.36	0.33	0.39	29.13	25.36	19.45	14.35	10.38	19.73	
T <sub>7</sub>	16.13	16.50	17.06	17.47	17.88	17.01	0.50	0.41	0.38	0.36	0.33	0.40	29.24	25.14	18.90	13.09	8.88	19.05	
T <sub>8</sub>	16.13	16.47	17.07	17.60	18.05	17.06	0.48	0.44	0.40	0.37	0.34	0.40	29.58	24.02	18.23	14.29	10.16	19.26	
Means	15.93	16.23	16.77	17.37	17.77	16.81	0.46	0.41	0.38	0.35	0.32	0.38	29.08	24.73	18.56	13.84	9.85	19.21	
			CD at 5%						CD at 5%								CD at 5%		
Treatment (A)			0.12						0.01								0.22		
Storage days (B)			0.09						0.010								0.17		
A × B			NS						NS								0.49		

for one hour using iced water at 5°C followed by air drying at room temperature. Each treatment was replicated thrice in Complete Randomized Block Design. After soaking, fruits were placed inside the polythene bags maintained at 2% level of ventilation and kept for observations under ambient conditions. Each treatment had fifty fruits per replication and sampled periodically at every one day interval. The fruits were assessed for physiological loss in weight, browning index, spoilage loss, TSS, titratable acidity and ascorbic acid content. For browning, the following scale was used (Ramma, 11): 0 = no browning (excellent quality); 1 = slight browning; 2 = 25% browning; 3 = 25–50% browning; 4 = 50–75% browning and 5 > 75% (very poor quality). Browning index was calculated as  $\bar{O}$  (browning scale x percentage of corresponding fruit within each class). Total soluble solids (TSS) was measured with help of hand refractometer. Titratable acidity and ascorbic acid content were evaluated according to the method described by Ranganna (12).

## RESULTS AND DISCUSSION

The fruit weight loss during the storage period of 4 days significantly increased with increasing period of storage. Lowest weight loss (2.57%) was noted in HWT (55°C, 12 min) i.e. T<sub>3</sub> which might be due to the recrystallisation or melting of the epicuticular wax, thereby reducing respiration and transpiration rate (Lichter *et al.*, 6). Whereas, treatment that involves a combination of HWT (98°C, 30 s) and OA (10%, 15 min) dipping in T<sub>8</sub> treatment was found to record the highest weight loss (5.08%) which might be due to a reduction in fruit firmness, indicating structural damage to the cross-linkages in the cell wall (Olesen *et al.*, 9) (Table 1).

Atinut *et al.* (1) reported that, higher the concentration of acid used greater is the degree of browning inhibition which supports the finding of higher browning index in fruit treated with HWT and OA at low concentration (0.25%). Maximum reduction (21.25) in browning was noted in T<sub>6</sub>, which was statistically *at par* with T<sub>8</sub>. This might be due to inhibition of PPO and POD activities. Highest browning index (377.50 and 445.83) in fruit treated with HWT (T<sub>3</sub> and T<sub>4</sub>) may possibly be due to extensive browning caused by increased anthocyanase activity as reported by Underhill and Critchley (16)

Percentage of spoilt fruits was found significantly highest in control i.e. T<sub>1</sub> (23.75%), which was statistically *at par* with T<sub>4</sub> and T<sub>2</sub>; while lowest spoilage (8.75%) in T<sub>6</sub> which was also statistically *at par* with T<sub>8</sub> (10.83) might be due to the fungistatic effects of the applied treatments - HWT by killing the organisms on and below the fruit

surface as reported by Fallik *et al.* (2) in apples, and OA by providing an acidic conditions on the peel surface that makes most fungi difficult to develop as confirmed by Lichter *et al.* (5) in litchi (Table 2).

Treatments showed a progressive increase in fruit TSS from the day just after treatment (16.12%) upto the end of the 4 days (17.97%) storage period. Significantly, maximum TSS (17.28%) was recorded in fruits treated with T<sub>6</sub> while lowest TSS (16.81%) was recorded in T<sub>1</sub>. Increase in TSS of fruit during storage at ambient storage conditions might be due to large loss of water from fruits (Ray *et al.*, 13) in which the concentration of sugars might have increased.

Titratable acidity of fruits was significantly affected by treatments and recorded lowest titratable acidity (0.35%) in T<sub>3</sub>, while highest titratable acidity (0.40%) was reported in T<sub>8</sub> and T<sub>7</sub> which was statistically *at par* with T<sub>6</sub>, T<sub>4</sub> and T<sub>1</sub>, whereas minimum acidity (0.35%) was recorded in T<sub>3</sub> which was also *at par* with T<sub>5</sub>. There occurred a gradual declining trend in acidity of fruit in all treatments with advancement of storage period, which might be due to utilization of organic acids in respiratory process and other biodegradable reactions (Mahajan, 7).

Throughout the entire storage period, massive reduction in ascorbic acid was observed regardless of treatments in which more than 50% loss was observed on the 3<sup>rd</sup> day of storage. Maintenance of higher ascorbic acid content ((19.73 mg/100g pulp) by T<sub>6</sub> treatment might be due to the inhibitory effect on ascorbic acid oxidation (Tannerbaum *et al.*, 15). Ascorbic acid being sensitive to light, oxygen and heat, its enormous reduction might be due to the fact that it was easily oxidize in the presence of oxygen by both enzymatic and non-enzymatic catalyst, and therefore, is liable to be lost during storage (Rai *et al.*, 10). Minimum ascorbic acid content (18.85%) was recorded in T<sub>2</sub> which was statistically *at par* with T<sub>1</sub>, T<sub>7</sub> and T<sub>4</sub> respectively

From the present study, it can be concluded that treatment with hot water (HWT) at 55°C for 12 min followed by dipping in oxalic acid (OA) at 10% for 15 min was the most effective treatment for maximum retention of physico-chemical parameters of litchi fruit at ambient temperature.

## REFERENCES

1. Atinut, J., Jammong, U., Kanda, W. and Kobkiat, S. 2007. Inhibition of pericarp browning after thawing of frozen litchi fruit by oxalic acid dipping. *J. Agril. Sci.* **38**: 45-48.
2. Fallik, E., Tuvia-Alkalai, S., Feng, X. and Lurie, S. 2001. Ripening characterization and decay

- development of stored apples after a short prestorage hot water rinsing and brushing. *Innovative Food Sci. Emerging Technol.* **2**: 127-32.
3. Fitzell, R.D. and Coates, L.M. 1995. Diseases of lychee. **In**: Coates, L., Cooke T., Persley, D., Beattie, B., Wade, N. and Ridgeway, R. (Eds.). *Postharvest Diseases of Horticultural Produce, Tropical Fruits*, Department of Primary Industries, Queensland, Australia, pp. 41-42.
  4. Kumar, A. and Kumar, G. 2004. Effect of foliar sprays of multi-K on yield quality and shelf life of litchi (*Litchi chinensis* Sonn.) cv. Rose Scented. *Advances-in-Plant-Sciences*, **17**: 519-523.
  5. Lichter, A.; Dvir, O.; Rot, I.; Akerman, M.; Feygenberg, O. and Pesis, E. 2004. Acidified peel of litchi fruits selects for postharvest *Penicillium* decay. *Phytoparasitica*, **32**: 226-36.
  6. Lichter, A., Dvir, O., Rot, I.; Akerman, M., Regev, R., Wiesblum, A., Fallik, E., Zauberman, G. and Fuchs, Y. 2000. Hot water brushing: an alternative method to SO<sub>2</sub> fumigation for colour retention of litchi fruits. *Postharvest Biol. Tech.* **18**: 235-44.
  7. Mahajan, B.V.C. 1997. Studies on the biochemical changes in litchi fruits during storage. *Indian J. Plant Physiol.* **2**: 310-11.
  8. Mirdehghan, S.H., Rahemi, M., Serrano, M., Guillen, F. and Martinez, D. 2006. Prestorage heat treatment to maintain nutritive and functional properties during postharvest cold storage of pomegranate. *J. Agric. Food Chem.* **54**: 8495-500.
  9. Olesen, T.; Wiltshire, N. and McConchie, C. 2003. Improved postharvest handling of lychee. *Rural Industries Research and Development Corporation Publication 03/111* pp.
  10. Rai, D.R.; Oberoi, H.S. and Baboo, B. 2002. Modified atmosphere packaging and its effect on quality and shelf life of fruit and vegetables- An overview. *J. Food Sci. Tech.* **39**: 199-207.
  11. Ramma, I. 2004. Post-harvest sulphur dioxide fumigation and low acid dip for pericarp color retention and decay prevention on litchi. *Revue-Agricole-et-Sucriere-de-l'Ile-Maurica*. **83**: 18-25.
  12. Ranganna, S. 1986. *Handbook of Analysis and Quality control for Fruit and Vegetable Products*. 2<sup>nd</sup> edn. Tata McGraw Hill Publishing Co. Ltd., New Delhi.
  13. Ray, P.K., Rani, R. and Singh, S.K. 2004. Effect of temperature and sulphur treatments on storage behavior of litchi fruits. *Indian J. Hort.* **61**: 292-95.
  14. Sivakumar, D., Regnier, T., Demoz, B. and Korsten, L. 2005. Effect of postharvest treatments on overall quality retention in litchi fruit during low temperature storage. *J. Hort. Sci. Biotechnol.* **80**: 32-38.
  15. Tannerrbaum, S.R., Young, V.R. and Archer, M.C. 1985. Vitamins and Minerals. **In**: Fennema, O.R. (Ed.), *Food Chemistry*. (2<sup>nd</sup> edn.). Marcel Dekker, Inc. New York. pp. 478-543.
  16. Underhill, S.J.R. and Critchley, C. 1993. Lychee pericarp browning caused by heat injury. *HortScience*, **28**: 721-22.
  17. Underhill, S.J.R. and Critchley, C. 1994. Anthocyanin decolourisation and its role in lychee pericarp browning. *Aust. J. Exp. Agric.* **34**: 115-22.
  18. Zauberman, G., Ronen, R., Akerman, M., Weksler, A., Rot, I. and Fuchs, Y. 1991. Postharvest retention of the red colour of litchi fruit pericarp. *Scientia Hort.* **47**: 89-97.
  19. Zheng, X., Tian, S., Gidley, M.J., Yue, H. and Li, B. 2007. Effects of exogenous oxalic acid on ripening and decay incidence in mango fruit during storage at room temperature. *Postharvest Biol. Tech.* **45**: 281-84.

---

Received: August, 2009; Revised: April, 2010

Accepted: August, 2010